

Ethnobotany of wild edible plants traditionally used by the local people in the Ramnagar region from Nainital District, Uttarakhand, India

Neetu Bohra^{*1}, Lalit M. Tewari² And Ashish Tewari³

¹⁻³ D.S.B. Campus, Kumaun University, Nainital

Email: neetubohra18@gmail.com

ABSTRACT

The present investigation deals with the ethnobotanical study conducted during the year 2014-2016 to collect, identify and record the wild edible plants (WEPs) used by the local as well as tribal people inhabiting in rural areas in the Uttarakhand state of ramnagar region of district Nainital, India. A total of 59 plant species belonging to 36 families are reported in this paper based on an ethnobotanical field study. The four major life forms were herbs, shrubs, climbers and trees. Trees (56%) made the highest proportion in this communication of the edible plant species followed by shrubs (25%), herbs (13%) and climbers (6%). Fruit, seed and leaves are found as most preferred edible plant parts. The plant species are arranged in alphabetical order by their botanical name with family, Habit, Altitudinal range (m), Flowering and Fruiting period, Local name, part/s used and ethnobotanical use.

Keywords: Wild edible plants, Ethnobotany, rural areas, Ramnagar, India.

INTRODUCTION

WEPs play an important role in ensuring food security and improve the nutrition in the diets of many people in developing countries (Lulekal et al., 2011; Ghorbani et al., 2012). They are potential sources of species for domestication and provide valuable genetic traits for developing new crops through breeding and selection (Ford-Lloyd et al., 2011).

The traditional knowledge on the use as plants is well documented, however, there is a

How to Site This Article:

Neetu Bohra, Lalit M. Tewari and Ashish Tewari (2017). Ethnobotany of wild edible plants traditionally used by the local people in the Ramnagar region from Nainital District, Uttarakhand, India. *Biolife*. 4(4), pp 12-19.

DOI: [10.5281/zenodo.7351167](https://doi.org/10.5281/zenodo.7351167)

Received: 2 January 2017;

Accepted; 19 February 2017;

Available online : 1 March 2017

lacuna in accurate documentation of the folk knowledge on WEPs. It is of outmost importance to document the folk knowledge on WEPs before this knowledge disappears. An inadequate understanding concerning taxonomy, biology and local knowledge of these plants has contributed to suboptimal utilization of these valuable resources. It is essential therefore to identify and evaluate these plants which can

broaden our choice of food and enrich the livelihoods of rural poor. The tradition of using wild palatable plants is still alive in the rural populations of Lesser Himalayas, but is vanishing. Consequently, the recording, preserving, and infusing of this traditional knowledge to upcoming generations is pressing and vital.


Still many more wild species are believed to be edible and undocumented yet. More recently, some ethnobotanical studies have undertaken in some parts of the country. However, the majority of these studies have the dealt with medicinal species and little emphasis has been paid to WEPs. This study has therefore sought to document indigenous knowledge related to uses of wild edible plant species and to assess the existing threats to WEPs in the study areas. Current over

harvesting of mature fruit from parent-tree influenced the regeneration of this species. If this unsustainable harvesting by local people continues, the capacity of the species to maintain its wild population is significantly reduced. Therefore, management and conservation strategies that incorporate this factor is required. Conservation and management of WEPs and fruits will help to enhance and maintain the regional biodiversity with minimal adverse impact on the biodiversity.

Although many ethnobotanical surveys have been conducted by different workers (Abbasi *et al.*, 2012; Hazarat *et al.*, 2011; Abbasi *et al.*, 2010; 2010; Abbasi *et al.*, 2009; & Hussain *et al.*, 2008) in different areas of Central Himalayas but to our knowledge no systematic investigation on ethno-botany of WEPs of Nainital district of Uttarakhand of central Himalayas, has been made. In this context, the present study will serve as a useful reference with particular emphasis on traditional uses of wild edible plants as there is dramatic loss of traditional knowledge regarding WEPs. The rapidity with which environmental damage, loss of floristic and cultural diversity occurs today, a necessity is felt for the recording and documentation of traditional knowledge about the uses of edible plants which is widely disappearing. Therefore, there is an urgent necessity to document traditional knowledge.

MATERIALS AND METHODS

The method employed in this study were designed with the purpose of providing baseline information on the use of plant species in local system, through literature survey and field visits to various areas from 2014-2016 in the Ramnagar region of Nainital district, Uttarakhand. The selected informants in the sample site were interviewed using semi-structured interview focusing on the WEPs. Full notes on facts and information about the respondents, history of wild food collectors, history of WEPs, and other essential information were recorded on site. A brief group discussion was also made with the informants. During the discussion the informants were free to state about WEPs and their knowledge without being interfered. Voucher specimen of each WEPs species was collected. Live specimens and photographs were shown to local villagers for local identification. The collected plants were identified with the help of specialist and available literature (Osmaston, 1927; Hooker,



RESULTS AND DISCUSSION

Photo Plate-1: Wild edible plants of Ramnagar region of Nainital district, Uttarakhand

Bauhinia variegata L.
(Kachnar)




Bombax ceiba L. (Semal)




Centella asiatica (L.)Urban
(Brahmi)



Collected tubers of *Costus speciosus* (Koeng ex Retz.)
Keula



Aegle marmelos Corr. (Bel)



Murraya koenigii (Linn.)Spreng.
(kari-pat)




Photo Plate 2: Wild edible plants of Ramnagar region of Nainital district, Uttarakhand

Phyllanthus emblica L.
(Aonla)




Solanum nigrum L. (Makoi)



Dioscorea bulbifera L.
(Gethi)



Colocasia esculenta (Linn.)Schott
(Pinalu)



Collected fruits of *Terminalia bellirica* Roxb. (Bahera)



Collected fruits of *Terminalia chebula* (Gaertn.) Retz. Harar



and infusing of this traditional knowledge to upcoming generations is pressing and vital.

Photo plate-3. Tribal woman busy in collecting *Chenopodium album* L. (Bathua) leaves for vegetable



Acknowledgement

Authors sincerely acknowledge the financial support provided by the IERP, GBPIHED, Kosi, Katarmal, Almora to carry out this research work, the informants who open-heartedly shared their knowledge with us and Prof Y.P.S. Pangtey for helping in identification of plant specimen.

Conflict of Interests

Authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1]. Abbasi, A.M., Khan, M.A., Ahmad, M., Zafar, M., Khan, H., Muhammad, N., Sultana, S.: Medicinal plants used for the treatment of jaundice and hepatitis based on socio-economic documentation. *Afric J Biotech* 2009, 8:1643-1650.
- [2]. Abbasi, A.M., Khan, M.A., Ahmad, M., Zafar, M., Jahan, S., Sultana, S.: Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North- West Frontier Province, Pakistan. *J Ethnopharmacol* 2010, 128:322-335.
- [3]. Abbasi, A.M., Khan, M.A., Ahmad, M., Zafar, M.: Medicinal plant biodiversity of Lesser Himalayas-Pakistan. New York: Springer; 2012:1-212.
- [4]. Arya, J.C.: Studies on medicinal flora of trai region of Uttarakhand. 2015, D.Phil submitted to D.S.B. Campus, Kumaun University Nainital.
- [5]. Badhwar, R. L., Fernandez, R.R.: Edible wild plants of the Himalayas. The forest Research Institute and Colleges, Dehradun
- [6]. Ford-Lloyd ,B.V., Schmidt, M., Armstrong, S.J., Barazani, O., Engels, J., Ge, S., Hadas, R., Hammer, K., Kell, S.P., Kang, D., Khoshbakht, K., Li, Y., Long, C.L., Lu, B.R., Ma, K.P., Nguyen, V.T., Qiu, L.J., Wei, W., Zhang, Z.W., Maxted, N.: Crop wild relatives-undervalued, underutilized and under threat? *BioScience*. 2011;9:559-565. doi: 10.1525/bio.2011.61.7.10. [Cross Ref]
- [7]. Gaur, R. D., 1999. Flora of Garhwal, Trans Media, Srinagar (Garhwal).
- [8]. Ghorbani, A., Langenberger, G., Sauerborn, J.: A comparison of the wild food plant use knowledge of ethnic minorities in Naban River Watershed National Nature Reserve, Yunnan, SW China. *J Ethnobiol Ethnomed*. 2012;9:17. doi: 10.1186/1746-4269-8-17. [PMC free article] [PubMed] [Cross Ref]
- [9]. Hazrat, A., Nisar, M., Shah, J., Ahmad, S.: Ethnobotanical study of some elite plants belonging to Dir, Kohistan valley, Khyber Pukhtunkhwa, Pakistan: *Pak J Bot* 2011, 43(2):787-795.
- [10]. Hooker, J. D.: Flora of British India. Vols. I - VI. L. Reeve and Company Ltd, The East House Book , Ashford, Kent, UK: (1875 -2006).
- [11]. Hussain, K., Shahzad, A., Hussnain, S.Z.: An Ethnobotanical survey of important wild medicinal plants of Hattar District Haripur, Pakistan, *Ethnobot Leaflets* 2008, 12:29-35.
- [12]. Khan, A., Mehmood, S., Khan, R.U., Khan, S.U.: Ecological, Taxonomical and Ethnobotanical Study of Wild Fruit Producing Xerophytes of District Bannu, Kpk. *Journal of Medicinal Plants Studies* 2013, Vol I (4): pp. 112-120
- [13]. Lulekal, E., Asfaw, Z., Kelbessa, E., Van, D., Amme P.: Wild edible plants in Ethiopia: a review on their potential to combat food insecurity. *Africa Focus*. 2011;9:71-121.
- [14]. Sateesh Pujari and Estari Mamidala (2015). Anti-diabetic activity of Physagulin-F isolated from *Physalis angulata* fruits. *The Ame J Sci & Med Res*, 2015,1(1):53-60
- [15]. Osmaston, A.E.: A forest flora for Kumaun. International Book Distributors, Dehradun, 1927, pp. 605.
- [16]. Tewari, L.M., Singh, N., Upreti, K., Pangtey, Y.P.S.: Medicinal plants of

Ranikhet. 2008, I Edition Consul Book Depot, Nainital

- [17]. Upreti, K., Tewari, L.M., Pangtey, Y.P.S., Jalal , J.S.,: diversity and distribution of wild edible fruit plants of Uttarakhand. Biodiversity potentials of the Himalaya, Gyanodaya Prakashan, Nainital 2010, pp. 157-196.